

BIBLIOGRAFÍA

- 1) Deepak M., Sheweta B. (2013). Composition, properties and health benefits of indigestible carbohydrate polymers as dietary fiber: A review. *International Journal of Biological Macromolecules*, 61, 1-6, [dx.doi.org/10.1016/j.ijbiomac.2013.06.044](https://doi.org/10.1016/j.ijbiomac.2013.06.044).
- 2) Kosseva M. R. (2013). Functional Food and Nutraceuticals Derived from Food Industry Wastes. En: Kosseva M. R., Webb C., Eds. *Food Industry Wastes. Assessment and Recuperation of Commodities* (pp. 103-119). EEUU: Academic Press. [dx.doi.org/10.1016/B978-0-12-391921-2.00006-8](https://doi.org/10.1016/B978-0-12-391921-2.00006-8).
- 3) Mussatto S. I., Dragone G., Roberto I. C. (2006). Brewer's spent grain: generation, characteristics and potential applications. *Journal of Cereal Science*, 43, 1-14, doi.org/10.1016/j.jcs.2005.06.001.
- 4) Steiner J., Procopio S., Becker T. (2015). Brewer's spent grain: source of value-added polysaccharides for the food industry in reference to the health claims. *European Food Research and Technology*, 241(3), 303-315, doi.org/10.1007/s00217-015-2461-7.
- 5) Mussatto S. I. (2009). Biotechnological Potential of Brewing Industry By-Products. En: *Biotechnology for Agro-Industrial Residues Utilisation*, P. Singh nee' Nigam, A. Pandey (Eds.), Springer, Dordrecht (pp. 313-326), doi.org/10.1007/978-1-4020-9942-7_16.
- 6) Freeman M., Gortner, R. A. (1932). The gums of cereal grains. *Cereal Chemistry*, 9, 505-518.
- 7) Fincher, G. B., & Stone, B.A. (1986). Cell walls and their components in cereal grain technology. En Y. Pomeranz (Ed.), *Advances in cereal science and technology* (pp. 207-295). St. Paul: American Association of Cereal Chemistry.
- 8) Fry S. C. (2004). Primary cell wall metabolism: tracking the careers of wall polymers in living plant cells. *New Phytologist*, 161(3), 641-675, doi.org/10.1111/j.1469-8137.2004.00980.x
- 9) Saeed, F., Pasha, I., Anjum, F. M., & Sultan, M. T. (2011). Arabinoxylans and arabinogalactans: A comprehensive treatise. *Critical Reviews in Food Science and Nutrition*, 51(5), 467-476, [dx.doi.org/10.1080/10408391003681418](https://doi.org/10.1080/10408391003681418).
- 10) Izydorczyk M. S., Dexter J.E. (2008). Barley β -glucans and arabinoxylans: Molecular structure, physicochemical properties, and uses in food products-a Review. *Food Research International*, 41, 850-868, doi.org/10.1016/j.foodres.2008.04.001.
- 11) Courtin, C. M., & Delcour, J. A. (1998). Physicochemical and breadmaking characteristics of low molecular weight wheat derived arabinoxylans. *Journal of Agricultural and Food Chemistry*, 46, 4066-4073, doi.org/10.1021/jf980339t.
- 12) Carvajal-Millan E., Guilbert S., Doublier J. -L., Micard V. (2006). Arabinoxylan/protein gels: structural, rheological and controlled release properties. *Food Hydrocolloids*, 20, 53-61, doi.org/10.1016/j.foodhyd.2005.02.011.
- 13) Lu Z. X., Walker K. Z., Muir J. G., O'Dea K. (2004). Arabinoxylan fibre improves metabolic control in people with type II diabetes. *European Journal of Clinical Nutrition*, 58(4), 621-628, [dx.doi.org/10.1038/sj.ejcn.1601857](https://doi.org/10.1038/sj.ejcn.1601857).
- 14) Cao L., Liu X., Qian T., Sun G., Guo Y., Chang F., Zhou S., Sun X. (2011). Antitumor and immunomodulatory activity of arabinoxylans: a major constituent of wheat bran. *International Journal of Biological Macromolecules*, 48, 160-164, doi.org/10.1016/j.ijbiomac.2010.10.014.
- 15) García A. L., Otto B., Reich S. C., Weickert M. O., Steiniger J., Machowetz A., Rudovich N. N., Möhlig M., Katz N., Speth M., Meuser F., Doerfer J., Zunft H. J., Pfeiffer A. H., Koebnick C. (2006). Arabinoxylan consumption decreases postprandial serum glucose, serum insulin and plasma total ghrelin response in subjects with impaired glucose tolerance. *European Journal of Clinical Nutrition*, 61, 334-341, doi.org/10.1038/sj.ejcn.1602525.

- 16) Vietor R. J., Angelino S. A. G. F., Voragen A. G. J. (1992). Structural features of arabinoxylans from barley and malt cell wall material. *Journal of Cereal Science*, 15(2), 213-222, [doi.org/10.1016/S0733-5210\(09\)80120-3](https://doi.org/10.1016/S0733-5210(09)80120-3).
- 17) Roberfroid M., Slavin J. (2000). Nondigestible oligosaccharides. *Critical Reviews in Food Science and Nutrition*, 40, 461-480, doi.org/10.1080/10408690091189239.
- 18) Vitali B., Ndagijimana M., Maccaferri S., Biagi E., Guerzoni M. E., Brigidi P. (2012). An *in vitro* evaluation of the effects of probiotics and prebiotics on the metabolic profile of human microbiota. *Anaerobe*, 18, 386-391, doi.org/10.1016/j.anaerobe.2012.04.014.
- 19) Roberfroid M. (2007). Prebiotics: the concept revisited. *The Journal of Nutrition Effects of Probiotics and Prebiotics*, 137, 831S-837S, doi.org/10.1093/jn/137.3.830S.
- 20) Gibson G. R., Hutkins R., Sanders M. E., Prescott S. L., Reimer R. A., Salminen S. J., Scott K., Stanton C., Swanson K. S., Cani P. D., Verbeke K., Reid G. (2017). The International Scientific Association for Probiotics and Prebiotics (ISAPP) consensus statement on the definition and scope of prebiotics. *Nature Reviews Gastroenterology & Hepatology*, 14, 491-502, doi.org/10.1038/nrgastro.2017.75.
- 21) Neyrinck A. M., Possemiers S., Druart C., Van de Wiele T., De Backer F., Cani P. D., Larondelle Y., Delzenne N. M. (2011). Prebiotic effects of wheat arabinoxylan related to the increase in bifidobacteria, Roseburia and Bacteroides/Prevotella in diet-induced obese mice. *PLoS One*, 6, e20944, doi.org/10.1371/journal.pone.0020944.
- 22) Grootaert C., Delcour J. A., Courtin C. M. Broekaert W. F., Verstraete W., Van de Wiele T. (2007). Microbial metabolism and prebiotic potency of arabinoxylan oligosaccharides in the human intestine. *Trends in Food Science & Technology*, 18, 64-71, doi.org/10.1016/j.tifs.2006.08.004.
- 23) Broekaert W. F., Courtin C. M., Verbeke K., VanDeWiele T., Verstraete W., Delcour, J. A. (2011). Prebiotic and other health related effects of cereal derived arabinoxylans, arabinoxylan-oligosaccharides, and xylooligosaccharides. *Critical Reviews in Food Science and Nutrition*, 51, 178-194, doi.org/10.1080/10408390903044768.
- 24) Zhou S., Liu X., Guo Y., Wang Q., Peng D., Cao, L. (2010). Comparison of the immunological activities of arabinoxylans from wheat bran with alkali and xylanase-aided extraction. *Carbohydrate Polymers*, 81, 784-789, doi.org/10.1016/j.carbpol.2010.03.040.
- 25) Zhang Z., Smith C., Li W. (2014). Extraction and modification technology of arabinoxylans from cereal by-products: A critical review. *Food Research International*, 65, 423-436, [dx.doi.org/10.1016/j.foodres.2014.05.068](https://doi.org/10.1016/j.foodres.2014.05.068).
- 26) Gruppen H., Hamer R. J., Voragen, A. G. J. (1991). Barium hydroxide as a tool to extract pure arabinoxylans from water-insoluble cell-wall material of wheat-flour. *Journal of Cereal Science*, 13, 275-290, [doi.org/10.1016/S0733-5210\(09\)80006-4](https://doi.org/10.1016/S0733-5210(09)80006-4).
- 27) Jacquemin L., Zeitoun R., Sablayrolles C., Pontalier P.-Y., Rigal L. (2012). Evaluation of the technical and environmental performances of extraction and purification processes of arabinoxylans from wheat straw and bran. *Process Biochemistry*, 47(3), 373-380, doi.org/10.1016/j.procbio.2011.10.025.
- 28) Wang J., Sun B., Liu Y., Zhang H. (2014). Optimisation of ultrasound-assisted enzymatic extraction of arabinoxylan from wheat bran. *Food Chemistry*, 150, 482-488, doi.org/10.1016/j.foodchem.2013.10.121.
- 29) Roos A. A., Persson T., Krawczyk H., Zacchi G., Stålbrand H. (2009). Extraction of water-soluble hemicelluloses from barley husks. *Bioresources Technology*, 100, 763-769, doi.org/10.1016/j.biortech.2008.07.022.
- 30) Fleury M., Edney M., Campbell L., Crow G. (1997). Total, water-soluble and acid-soluble arabinoxylans in western Canadian barleys. *Canadian Journal of Plant Science*, 77(2), 191-196, doi.org/10.4141/P96-033.

- 31) Ganguli N. K., Turner M. A. (2008). A simplified method for extracting water-extractable arabinoxylans from wheat flour. *Journal of the Science of Food and Agriculture*, 88 (11), 1905-1910. doi.org/10.1002/jsfa.3273.
- 32) Fadel A., Ayman M., Mahmoud A. M., Ashworth J. J., Li W., Ng Y. L., Plunkett A. (2018). Health-related effects and improving extractability of cereal arabinoxylans. *International Journal of Biological Macromolecules*, 109, 819-831, doi.org/10.1016/j.ijbiomac.2017.11.055.
- 33) Garrote G., Domínguez H., Parajó J. (1999). Hydrothermal processing of lignocellulosic materials. *European Journal of Wood and Wood Products*, 57(3), 191-202, doi.org/10.1007/s001070050039.
- 34) Doner L. W., Hicks, K. B. (1997). Isolation of hemicellulose from corn fiber by alkaline hydrogen peroxide extraction. *Cereal Chemistry*, 74, 176-181, dx.doi.org/10.1094/CCHEM.1997.74.2.176.
- 35) Pan, G. X., Bolton, J. L., Leary, G. J. (1998). Determination of ferulic and *p*-coumaric acids in wheat straw and the amounts released by mild acid and alkaline peroxide treatment. *Journal of Agricultural and Food Chemistry*, 46, 5283-5288, doi.org/10.1021/jf980608f.
- 36) Höjje A., Gröndahl M., Tømmerraas K., Gatenholm P. (2005). Isolation and characterization of physicochemical and material properties of arabinoxylans from barley husks. *Carbohydrate Polymers*, 61, 266-275, doi.org/10.1016/j.carbpol.2005.02.009
- 37) Peng F., Peng P., Xu F., Sun R. C. (2012). Fractional purification and bioconversion of hemicelluloses. *Biotechnology Advances*, 30, 879-903, doi.org/10.1016/j.biotechadv.2012.01.018.
- 38) Xu, F., Liu, C. F., Geng, Z. C., Sun, J. X., Sun, R. C., Hei, B. H., Lin L., Wu S. B. Je J. (2006). Characterisation of degraded organosolv hemicelluloses from wheat straw. *Polymer Degradation and Stability*, 91(8), 1880-1886, doi.org/10.1016/j.polymdegradstab.2005.11.002.
- 39) Escarnot E., Aguedo M., Paquot M. (2012). Enzymatic hydrolysis of arabinoxylans from spelt bran and hull. *Journal of Cereal Science*, 55, 243-253, doi.org/10.1016/j.jcs.2011.12.009.
- 40) Li W., Hu H., Wang Q., Brennan, C. S. (2013). Molecular features of wheat endosperm arabinoxylan inclusion in functional bread. *Foods*, 2(2), 225-237, doi.org/10.3390/foods2020225.
- 41) Alonso J. L., Domínguez H. X. P. A. P. T. Garrote G., Parajó J., Vázquez M. J. (2003). Xylooligosaccharides: Properties and production technologies. *Electronic Journal of Environmental, Agricultural and Food Chemistry*, 2, 230-232.
- 42) Pitkanen L., Tuomainen P., Virkki L., Tenkanen M. (2011). Molecular characterization and solution properties of enzymatically tailored arabinoxylans. *International Journal of Biological Macromolecules*, 49(5), 963-969, dx.doi.org/10.1016/j.ijbiomac.2011.08.020.
- 43) Ebringerova A., Hromadkova Z. (2002). Effect of ultrasound on the extractability of corn bran hemicelluloses. *Ultrasonics Sonochemistry*, 9(4), 225-229, doi.org/10.1016/S1350-4177(01)00124-9.
- 44) Hromádková Z., Kováčiková J., Ebringerová A. (1999). Study of the classical and ultrasound-assisted extraction of the corn xylan. *Industrial crop and products*, 9, 101-109, doi.org/10.1016/S0926-6690(98)00020-X.
- 45) Roos A. A., Persson T., Krawczyk H., Zacchi G., Ståhlbrand H. (2009). Extraction of water-soluble hemicelluloses from barley husks, *Bioresource Technology*, 100 (2), 763-769, doi.org/10.1016/j.biortech.2008.07.022.
- 46) Official Methods of Analysis of AOAC International, Solids (total) and Moisture in Flour - Air Oven Methods (32.1.03). Chapter 32(1). Maryland: USA, 2006.

- 47) AOAC 942.05. 2000. Ash of animal feed. Gaithersburg, MD, USA.
- 48) IUPAC. (1992). Standard Methods for the Analysis of oils, Fats and Derivates. 7th ed. Eds. Paquot C, Hautffenne A. International Union of Pure and Applied Chemistry, Blackwell Scientific Publications Inc., Oxford, UK.
- 49) https://www.megazyme.com/documents/Booklet/K-TDFR-200A_DATA.pdf
- 50) Dubois, M., Gilles, K. A., Hamilton, J. K., Rebers, P. A., Smith, F. (1956). Colorimetric method for determination of sugars and related substances. *Analytical Chemistry*, 28(3), 350-356, doi.org/10.1021/ac60111a017.
- 51) Blumenkrantz, N., Asboe-Hansen, G. (1973). New method for quantitative determination of uronic acids. *Analytical Biochemistry*, 54(2), 484-489, doi.org/10.1016/0003-2697(73)90377-1.
- 52) Carrier M., Loppinet Serani A., Denux D., Lasnier J., Ham Pichavant F., Cansell F., Aymonier C. (2011). Thermogravimetric analysis as a new method to determine the lignocellulosic composition of biomass. *Biomass Bioenergy*, 35(1), 298-307. doi.org/10.1016/j.biombioe.2010.08.067.
- 53) Glasser W. G., Kaar W. E., Jain R. K., Sealey, J. E. (2000). Isolation options for non-cellulosic heteropolysaccharides (HetPS). *Cellulose*, 7, 299-317, doi.org/10.1023/A:1009277009836.
- 54) Huebner J., Wehling R.L., Hutkins R.W. (2007). Functional activity of comercial prebiotics. *International Dairy Journal*, 17, 770-775, doi.org/10.1016/j.idairyj.2006.10.006.
- 55) Reid G., Kim S.O., Köhler G.A. (2006). Selecting, testing and understanding probiotic microorganisms. *FEMS Immunology and Medical Microbiology*, 46, 149-157, doi.org/10.1111/j.1574-695X.2005.00026.x.
- 56) Moongngarm A., Trachoo N., Sirigungwan N. (2011). Low molecular weight carbohydrates, prebiotic content, and prebiotic activity of selected food plants in Thailand. *Advance Journal of Food Science and Technology*, 3, 269-274.
- 57) Gómez E., Tuohy K.M., Gibson G.R., Klinder A., Costabile A. (2010). *In vitro* evaluation of the fermentation properties and potential prebiotic activity of Agave fructans. *Journal of Applied Microbiology*, 108, 2114-2121, doi.org/10.1111/j.1365-2672.2009.04617.x.
- 58) Rubel I. A., Pérez E. E., Genovese D.B., Manrique G.D. (2014). *In vitro* prebiotic activity of inulin-rich carbohydrates extracted from Jerusalem artichoke (*Helianthus tuberosus* L.) tubers at different storage times by *Lactobacillus paracasei*. *Food Research International*, 62, 59-65, dx.doi.org/10.1016/j.foodres.2014.02.024.
- 59) Martínez-Villaluenga C., Gómez R. (2007). Characterization of bifidobacteria as starters in fermented milk containing raffinose family of oligosaccharides from lupin as prebiotic. *International Dairy Journal*, 17, 116-122, doi.org/ 10.1016/j.idairyj.2006.02.003.
- 60) Jouany J. P. (1982). Volatile fatty acids and alcohol determination in digestive contents, silage juice, bacterial cultures and anaerobic fermentation contents. *Science des Aliments*, 2, 131-144.